

The Hazard Evaluation System and Information Service: A Physician's Resource in Toxicology and Occupational Medicine

KIM HOOPER, PhD, Berkeley, California

Hazard evaluation is an emerging science. The Hazard Evaluation System and Information Service (HESIS), part of California's program in preventive occupational health, is a resource for clinicians who wish to stay abreast of the relationship between toxicology and occupational health. For example, advances in assays for cancer or reproductive effects in test animals enable us to identify with greater confidence significant cancer or reproductive hazards among the increasing variety of workplace exposures. Occupational experiences with dibromochloropropane (DBCP), Kepone, bis(chloromethyl) ether, benzidine and vinyl chloride demonstrate the shortcomings of relying on human data. The latency period of cancer, limited sensitivity of epidemiologic studies and severity of effects require us to use animal test data to evaluate the potential cancer and reproductive risks of workplace substances. HESIS gives appropriate weight to experimental data in hazard evaluations of chemicals such as ethylene oxide, ethylene dibromide, polychlorinated biphenyls and the glycol ethers. A similar approach is apparent in the California Department of Health Services' recently released Carcinogen Identification Policy.

In 1977 male workers engaged in formulating commercial batches of several pesticides, including the nematocide dibromochloropropane (DBCP), at a chemical plant in Lathrop, California, realized that for several years none of them had fathered a child. Their fears of apparent infertility were confirmed when, with the cooperation and assistance of their union, the company and health professionals, semen analyses were conducted and the group was found to have an abnormally high prevalence of oligospermia (low sperm count) and azospermia (absence of sperm).¹ The frequency and severity of these effects correlated well with the duration of the men's exposure to DBCP. Pronounced effects on spermatogenesis occurred at exposure levels as low as 0.4 ppm (eight-hour time-weighted average), doses that produced no other clinical signs of toxicity.²

Shortly thereafter, results from the National Cancer Institute Bioassay program were released showing that DBCP caused cancer in test animals at doses very close to those that workers at Lathrop might have received.³ In addition to being a potent carcinogen, DBCP was also shown to be genotoxic in several short-term tests for mutagenicity using bacteria⁴⁻⁶ or mammalian cells.^{7,8}

The DBCP incident was the first documented example of workplace-induced reproductive failure in men. The rapidly differentiating reproductive tissue in male test animals seems to be particularly sensitive to certain chemicals (see Ethylene Dibromide later in this paper) as is the differentiating fetal tissue in pregnant women (see Glycol Ethers). Public concern was heightened by the discovery that studies were published 18 years earlier showing that DBCP caused testicular

Refer to: Hooper K: The Hazard Evaluation System and Information Service: A physician's resource in toxicology and occupational medicine, *In Occupational disease—New vistas for medicine*. West J Med 1982 Dec; 137:560-571.

Dr. Hooper is Chief, Hazard Evaluation System and Information Service, State of California Department of Health Services and Department of Industrial Relations, Berkeley.

Reprint requests to: Kim Hooper, PhD, Chief, Hazard Evaluation System and Information Service, State of California Department of Health Services and Department of Industrial Relations, 2151 Berkeley Way, Berkeley, CA 94704.

ABBREVIATIONS USED IN TEXT

DBCP=dibromochloropropane
 HESIS=Hazard Evaluation System and Information Service
 TIRS=Telephone Inquiry Response System

changes in test animals at dose levels close to those that produced similar adverse spermatogenic effects in the men.⁹ Had greater significance been attached to these results in test animals, the reproductive failure in workers at Lathrop might have been avoided.

What can be done to help prevent similar episodes from occurring in the future? Two lessons may be learned. First, the release of information at the right time (for example, animal toxicity data on DBCP to plant officials and workers at Lathrop) could be an effective part of a program of preventative occupational medicine. Second, greater weight may need to be given to results from studies in test animals as predictors of effects in humans.

Shortly after the DBCP episode, the California legislature moved to strengthen the state's academic resources in occupational health and medicine by establishing several Occupational Health Centers on campuses of the University of California. Right-to-know legislation in California was enacted to enable employees and employers to receive information on the hazards of workplace substances and to permit employees (or their physicians) to have access to personal medical records. Finally, the legislature created the Hazard Evaluation System and Information Service (HESIS) within the State's Department of Health Services.*

Hazard Evaluation System and Information Service

HESIS practices preventive occupational medicine by providing information to workers and employers on the health effects of toxic substances used in workplaces and by assisting alert and concerned clinicians in identifying possible occupational determinants of disease. The HESIS scientific staff of occupational health physicians and PhD toxicologists attempts to bridge the gap between the published scientific and medical literature and the workplace community (employers, employees and health professionals) by a systematic review of current issues of 57 journals in toxicology, epidemiology, occupational medicine and related fields. These resources are present in the HESIS library. To provide the most up-to-date information possible on questions of toxicology and occupational health, the HESIS computer provides access to bibliographies of the world's published medical and toxicologic literature, US-sponsored research in progress and numerous medical, toxicologic and occupational health data bases.

HESIS provides information to the occupational health community in several ways. A Telephone Inquiry

Response System (TIRS) is available to answer questions from employees, employers or health professionals about clinical symptoms or about toxic substances encountered in California's workplaces. HESIS staff, including scientists, health educators and a librarian, responded to over 1,000 such inquiries in 1981. Such calls may alert the medical community to new health hazards or may trigger follow-up investigations or remedial consultations.

TIRS serves as a listening post for the health problems of California's workplaces. With TIRS and a continual review of the scientific and medical literature and of data from various health registries (such as data on tumors and birth defects), HESIS has begun to create a reasonably comprehensive occupational health surveillance system for California. The cooperation of physicians is needed to improve and extend this system.

From such surveillance, HESIS on occasion conducts an in-depth investigation of a targeted substance, group of substances, health effect or occupation, and releases a written review of the scientific and medical evidence that documents the potential for hazard.¹⁰⁻²² Following this evaluation, HESIS may inform the workplace community of any new or unappreciated health hazard by issuing a *Hazard Alert*,²³⁻²⁵ *Information Bulletin*²⁶ or *Fact Sheet for Physicians*,²⁷⁻³³ depending upon the apparent severity of the problem. As a consequence of such a hazard evaluation, HESIS may also recommend to Cal-OSHA that the work practices or permissible exposure level (PEL) for a substance be changed (examples are ethylene oxide [EtO],²³ glycol ethers²⁴ and ethylene dibromide [EDB].²⁵)

Cancer and Reproductive Hazards in the Industrial Chemical World

Over the past 40 years the production of synthetic organic chemicals has increased enormously (greater than 300-fold),³⁴ and workers are exposed to a much greater variety of chemicals than ever before. Few of these approximately 50,000 substances have been tested for their ability to cause such severe and life-threatening effects as cancer, mutation, birth defects or sperm damage. For those that have been studied, it has not been clear how animal test data, which provide most of the evidence, should best be used.

The long latency period of some human cancer (as long as 20 to 40 years)³⁵ raises the possibility that past and present workplace exposures will cause increases in cancer among workers in the future. Although the cancer rate of the general population at present appears to be relatively stable, lung cancer has increased significantly over the last 80 years.³⁶ The major portion of this increase is attributed to increased cigarette smoking,^{37,38} but the contributions of occupational factors are less clear.^{37,39} What is clear is that significant increases in cancer have occurred in certain workplaces during this same period—such as exposures to bis(chloromethyl) ether, asbestos, 2-naphthylamine and benzidine—without causing visible changes in the cancer incidence in the general population, and that unacceptably high cancer risks were experienced by indi-

*HESIS is in the Toxic Substances Control Division of the state's Department of Health Services, 2151 Berkeley Way, Berkeley, CA 94704 (415/540-3014), and is funded by a contract with Cal/OSHA in the Department of Industrial Relations.

HAZARD EVALUATION SYSTEM

vidual workers from their exposure to carcinogenic agents.

Most of the known human carcinogens to which the public may be exposed have been identified by the high cancer rates observed among workers in specific industries: of the 36 agents for which there is strong or conclusive epidemiologic evidence of carcinogenic effect in humans, 26 were identified by occupational exposures; the remainder are mainly agents used in cancer chemotherapy (see Table 1).⁴⁰

Although less time may be required to see the effects in workers from exposures to mutagenic or reproductive toxins, very little is known about the prevalence of these effects in workplace situations.⁴¹ Effects on spermatogenesis have been studied for occupational exposures to only eight chemicals; clear or suggestive evidence is present for four chemicals other than DBCP^{1,42-44} (lead,⁴⁵ carbon disulfide,^{46,47} carbaryl^{48,49} and dinitrotoluene/toluenediamine⁵⁰); and no effects were seen with three others: polybrominated biphenyl (PBB),⁵¹ p-tertiary butylbenzoic acid⁵² and epichlorohydrin.⁴² Adverse reproductive effects in male and fe-

male workers have been associated with workplace exposures to several substances.⁵³ In some cases, spontaneous abortions may arise from damage to sperm cells (see Table 2).

Cancer and reproductive disabilities are serious health hazards. They compel us to regulate workers' exposures to toxic substances and to use animal data where possible to prevent such outcomes from occurring in humans.

Assessment of Reproductive Hazards:

The Value of Animal Data

The DBCP incident raises an important question: How can test results from studies of substances in ex-

TABLE 1.—Chemicals, Groups of Chemicals and Industrial Processes That Are Carcinogenic or Probably Carcinogenic to Humans

Carcinogenic

4-Aminobiphenyl
Arsenic and certain arsenic compounds
Asbestos
Manufacture of auramine
Benzene
Benzidine
N,N-bis(2-chloroethyl)-2-naphthylamine (chlornaphazine)*
Bis(chloromethyl)ether and technical grade chloromethyl methyl ether
Chromium and certain chromium compounds
Diethylstilbestrol*
Underground haematite mining
Manufacture of isopropyl alcohol by the strong acid process
Melphalan*
Mustard gas
2-Naphthylamine
Nickel refining
Soots, tars and mineral oils
Vinyl chloride

Probably Carcinogenic

Acrylonitrile
Aflatoxin*
Amitrole (aminotriazole)
Auramine
Beryllium and certain beryllium compounds
Cadmium and certain cadmium compounds
Carbon tetrachloride
Chlorambucil*
Cyclophosphamide*
Dimethylcarbamoyl chloride
Dimethylsulphate
Ethylene oxide
Iron dextran*
Nickel and certain nickel compounds
Oxymetholone*
Phenacetin*
Polychlorinated biphenyls
Tris(1-aziridinyl)phosphine sulphide (thiotepa)

*Evidence for effects in humans was obtained from nonoccupational exposures. Adapted from IARC, 1979.⁴⁰

TABLE 2.—Adverse Reproductive Effects Reportedly Associated With Occupational Exposures*

Chemical	Reported Effects
<i>To Women</i>	
Laboratory solvents ⁵⁴⁻⁵⁶ ...	(Chromosome aberrations); (birth defects)
Lead and/or other smelter exposures ⁵⁷⁻⁶⁰ ..	Spontaneous abortions; reduced birth weights; (congenital abnormalities)
Mercury (metal) ⁶¹	(Abnormal ovarian function)
Anesthetic gases ⁶²⁻⁷⁰	Spontaneous abortions; reduced birth weights; congenital malformations
Phthalate esters ⁷¹	(Anovulation); (increased spontaneous abortions)
Formaldehyde ⁷²	Menstrual disorders; (increased spontaneous abortions); reduced birth weights
Carbon disulfide ^{73,74}	Menstrual disorders; increased spontaneous abortions; (reduced fertility)
Organochlorine pesticides ⁷⁵⁻⁷⁷	(Abnormal ovarian function); (increased toxemia of pregnancy); (impaired lactation)
Benzene, toluene xylene ⁷⁸ .	Prolonged menstrual bleeding
Pesticides (various) ⁷⁹⁻⁸³ ...	Chromosome aberrations
<i>To Men</i>	
Dibromochloropropane (DBCP) ^{1,2,45,84,85}	Infertility, azoospermia, oligospermia
Lead and/or other smelter exposures ^{45,57,86-88}	Spontaneous abortions; premature births; [chromosomal abnormalities]; sperm abnormalities
Vinyl chloride ⁸⁹⁻⁹⁵	[Chromosomal abnormalities]
Anesthetic gases ^{95,96,98}	Increased spontaneous abortions; congenital abnormalities
Kepone ⁹⁶	Loss of libido; reduced sperm counts
Carbon disulfide ⁴⁷	Decreased libido; impotence; increased sperm abnormalities
Ethylene dibromide ⁹⁷	(Decreased fertility)

Evidence is weaker for reported effects in ().
Events reported in nonreproductive tissues in [].

*Adapted from Council on Environmental Quality.⁵³

HAZARD EVALUATION SYSTEM

TABLE 3.—Comparison of Reported Teratogenic Effects in Humans and Experimental Animals for Seven Agents*

Agent	Reported Sites in Humans	Reported Sites in Animals
Anesthetic gases	Hemangiomas, hernias, skin, heart	Skeletal defects only: rat, mouse
Smelter emissions (lead and/or arsenic)	Multiple malformations	Multiple malformations: rat, mouse, hamster (lead and arsenic)
Polychlorinated biphenyls	Skin discoloration; enlarged fontanelles	Skin discoloration and lesions: rhesus monkey; enlarged fontanelles and syndactyly: pig, dog
Alcohol	Facial, central nervous system	Facial, dermal, neural, extremities: rat, mouse
Vinyl chloride	(Neural tube)	Various, including encephalocele: rat
Diphenylhydantoin	Cleft lip, cleft palate, other craniofacial, mental deficiency	Cleft lip, cleft palate, syndactyly, other skeletal defects: mouse; minor kidney anomalies: rhesus monkey
Methylmercury	Central nervous system	Central nervous system, skeletal: rat, mouse, hamster, cat

Evidence is weaker for reported effects in ().

*Adapted from Council on Environmental Quality.⁵³

TABLE 4.—Comparison of Lowest Effective Doses of Eight Teratogens in Humans and Animals*

Chemical	Species	Ratio of Animal Dose: Human Dose
Thalidomide ^{102,103}	Rabbit	5.0-2.5
Polychlorinated biphenyls ^{104,105}	Rhesus monkey	1.8
	Dog	14.3
Alcohol ^{106,107}	Rat	3.8-7.6
Aminopterin ^{108,109}	Rat	2.0
Methotrexate ^{110,111}	Rat	4.8
Methylmercury ¹¹²⁻¹¹⁴	Cat, rat	50.0
Diethylstilbestrol (DES) ^{115,116}	Rhesus monkey	10.0-2.5
Diphenylhydantoin ^{117,118}	Mouse	25.0

*Adapted from Council on Environmental Quality.⁵³

perimental animals best be evaluated and communicated so as to protect worker health? The interspecies comparisons that can be made between humans and animals for reproductive endpoints (such as spermatotoxicity and teratogenicity) are interesting but are based on very limited data. Comparisons that have been made for semen parameters (counts, motility or morphology) between at least two species (including man) show a reasonably similar response for 24 chemicals.⁹⁸ Epidemiologic and laboratory studies that assay for semen quality (counts, morphology, or motility) in humans or animals are relatively easy to conduct and quite sensitive. If appropriate human study populations can be identified, a more meaningful comparison between the spermatotoxic effects of chemicals in animals and humans may be made in the near future.

Teratogenicity

For teratogenic or embryotoxic endpoints, several thousand substances have been tested in experimental animals, and several hundred have been found to give a positive result.⁹⁹ Only a handful have been studied in humans, and fewer of these produce effects.¹⁰⁰ The infrequency of teratogenic events limits the sensitivity of epidemiologic studies, and makes it unlikely that the number of identified human teratogens will increase substantially in the near future.¹⁰¹ Thus, our knowledge of the predictive value of animal teratogenicity data will likely remain limited. Nevertheless, for the several

agents that have been examined, comparisons of their teratogenic effects and potencies reveals that the human and animal responses are not as dissimilar as might have been expected (see Tables 3 and 4). Thus, for reproductive toxins in general, there may be more of an overlap than we had imagined between agents that affect humans and those that produce effects in test animals.

Cancer Prevention in the Workplace

A first step in limiting cancer risks to individuals and the working population is to reduce the causes of major varieties of cancer—lung, colon-rectal and breast—and control exposures to specific cancer-causing substances. The major identified causes of lung cancer include tobacco smoke and asbestos,^{38,119} and diet is believed to play an important role in breast and colon-rectal cancer.¹²⁰ In theory, cancer may be prevented by modifying the diet, controlling the use of tobacco and reducing exposures to cancer-causing substances such as asbestos. As a practical matter, involuntary environmental or occupational exposures may be easier to control than has been the voluntary use of tobacco or improper diet.

While attention has been focused on assessing the relative proportion of present and future cancer incidences that are caused by either "lifestyle" or occupational and environmental factors^{37,39} (with varying estimates that need to be resolved by future studies³⁹), an uncertainty of perhaps greater public health consequence may be overlooked. The effect of simultaneous exposures of workers to both factors, as is likely to occur in the workplace and in everyday life, is more relevant to an appropriate assessment of human risk. In two well-documented human studies of cancer risks from exposures to agents singly or in combination (cigarette smoking and exposures to asbestos or radiation), the cancer risks from the combined exposures are the product, not the sum, of the risks from separate exposures. Thus, prolonged cigarette smoking is associated with a tenfold increase in the risk for bronchial carcinoma (lung cancer) while prolonged heavy occupational exposure to asbestos is associated with a fivefold increase in risk. However, combined exposures

(such as cigarette smokers exposed to asbestos) are associated with a 50-fold, and not 15-fold, increase in risk over that experienced by nonsmokers with no asbestos exposure.¹¹⁹ There are data to suggest that a similar synergism exists between exposures to tobacco smoke and radiation.

Such synergistic interactions may be more common than is currently appreciated, but at present we cannot predict whether the effects of other interactions will be additive, multiplicative or inhibitory. Thus, from a policy as well as a scientific standpoint, the dichotomy between "lifestyle" and environmental and occupational factors may be more artificial than helpful. A simpler view is that cancer has a multiplicity of interacting causes, including "lifestyle" and environmental and occupational exposures, and that these will only infrequently be disentangled.

Oncogenes and Cancer

Oncogenes¹⁴⁹⁻¹⁵⁶ have been isolated from tumors of several human tissues (bladder, breast and lung), the sites of the major types of cancer in humans. When introduced into normal cells, these genes transform the cells into a cancerous state. Genes with similar transforming abilities have been isolated from tumors of these same tissues from test animals. Whereas distinctly different cancer genes exist for bladder, breast and lung, it is significant that the oncogenes from human and test animals are virtually identical for a specific tissue. In addition, some of these genes are virtually identical to the cancer-causing genes of the common tumor viruses that produce cancers in animals.

Even more remarkable is the finding that such genes are present in normal cells where they may participate in the maintenance of daily cellular functions. Damage to these genes (or to other controlling genes) by carcinogens or tumor virus infection may stimulate their activity and lead to cancer.

While much remains to be described of the mechanism of action of cancer genes, the recent discoveries suggest a unifying theory of cancer-causation that is gratifyingly simple. Normal cells contain certain genes necessary for normal cellular function. Modification (mutation, rearrangement and the like) of these genes (or of other genes that normally restrain their activity) by carcinogens or tumor virus infection causes them to function abnormally. This genetic misbehavior converts the cell to cancerous growth.

Methods for Identifying Carcinogens

A carcinogen is generally understood to be a substance or agent that increases the frequency (age-specific incidence) of cancer in humans or in other species.^{40,121-129} The identification of chemical substances that pose cancer risks to humans is complex and requires integration of information from several scientific disciplines. Evidence that a substance may be a carcinogen comes from four sources: epidemiologic studies in human populations; bioassays in experimental animals; short-term tests for mutagenicity and cell transformation, and similarities in chemical struc-

ture to known carcinogens. Procedures for evaluating data from the first three of these methods are reviewed briefly; it should be noted that they provide evidence of different strengths.

Epidemiologic Studies

Epidemiologic studies offer the overwhelming advantage of providing direct evidence for carcinogenic effects in *humans*. They are well suited to identify major causes of cancer in defined populations (cigarettes, asbestos, initial pregnancy late in life and so forth)¹³⁰ but are less suited to determining whether a *specific* chemical poses a cancer risk to humans. Even so, epidemiology has identified several important chemical carcinogens (such as benzene, arsenic) before animal tests were done.⁴⁰ At present, the International Agency for Research on Cancer states that there is sufficient evidence for the carcinogenicity in humans of 18 chemicals, groups of chemicals and industrial processes, and there is probable evidence of varying strengths for 18 others⁴⁰ (see Table 1).

In addition, negative results from well-conducted epidemiologic studies are useful in placing an upper limit of risk for a chemical exposure. They complement the more uncertain quantitative risk estimates made from data from animal cancer tests.

Epidemiologic methods are extremely useful tools. In general, however, they have low sensitivity. For example, when a cohort study population is relatively small (less than 1,000), the study may fail to identify an agent that increases the risk of a specific type of cancer by a factor of less than five to ten. Even large-scale studies may require an increase of more than 50 percent in cancer incidence before an effect is statistically significant. For this reason, negative results in studies of smaller size seldom provide strong evidence that an agent is not carcinogenic. To detect a carcinogenic effect, it may be necessary to have either a very large study population or a smaller-sized population that is exposed for several years to large doses of a potent carcinogen. In addition, such studies are often difficult to conduct, both because appropriate study groups and reliable information about past exposures are limited and because biases and confounding factors are difficult to eliminate. A recent review¹³¹ reports that epidemiologic data do not exist (and are unlikely to be developed in the future) for the vast majority of industrial chemicals that cause cancer in experimental test animals and to which workers are exposed.

In addition, because of the 20- to 30-year latency period of many types of cancer in humans, epidemiologic studies are not suited to warn and protect people from the cancer risks from exposures to new carcinogens. If an early-stage carcinogen has been identified by an epidemiologic study as a cause of human cancer and the exposures are reduced or eliminated, the cancer risk among those previously exposed may remain appreciable for the ensuing 20 to 30 years. Limiting exposure to late-stage carcinogens or promoters can reduce the cancer risk more rapidly—for example, ces-

sation of cigarette smoking appears to modify the risk for lung cancer within five years.^{132,133}

Thus, because of the insensitivity of epidemiologic studies, the long latency period of cancer and the difficulty in obtaining an appropriate study population, we are forced to rely heavily on other means of identifying agents which have the potential to produce cancer in humans.

Assessment of Cancer Hazards: The Value of Animal Data

Fortunately, results from animal cancer bioassays appear to be reasonable qualitative predictors of carcinogenic effects in humans, and the laboratory animal bioassay is widely used to indicate the carcinogenic potential of a chemical. Bioassay methods have been standardized in recent years and, except for minor details, there is now general acceptance of test procedures.^{125,134-140} Most substances that are carcinogenic in one species of test animal are carcinogenic in a second when adequately tested;¹⁴¹⁻¹⁴⁸ most substances that are known to be carcinogenic in humans, for which adequate animal data exist, are carcinogenic to animals.^{40,146-148} For several recognized human carcinogens—4-aminobiphenyl, bis(chloromethyl) ether, diethylstilbestrol, melphalan, mustard gas and vinyl chloride—the first evidence of carcinogenicity was found in test animals. Only afterwards were cancer effects looked for, and found, in humans.¹⁴⁸ From a scientific standpoint, it seems reasonable to consider substances for which there is evidence of toxic effects in test animals as likely to produce similar effects in humans. Thus, the International Agency for Research on Cancer concludes, "In the absence of adequate data in humans it is reasonable, for practical purposes, to regard chemicals for which there is *sufficient evidence* of carcinogenicity (i.e., a causal association) in animals *as if they presented a carcinogenic risk for humans*"⁴⁰ [emphasis added].

Indeed, recent discoveries in the molecular biology of cancer would make it surprising if carcinogenesis in rodents is greatly different from that in humans. Cancer-causing genes ("oncogenes") and cell transformation maintenance factors have been isolated from rodent tissues and are virtually identical to those found in the corresponding human tissues.¹⁴⁹⁻¹⁵⁶ These findings indicate that carcinogenesis in humans and test animals may be remarkably similar, and they strongly support the belief that test animals are reasonable and appropriate models for understanding the carcinogenic process in humans.

Sufficient evidence presently exists for the carcinogenicity in animals of about 200 chemicals. The International Agency for Research on Cancer considers there is "sufficient evidence" for carcinogenicity for 142 of the 422 chemicals it has assessed in its review process.⁴⁰ The National Cancer Institute has concluded that there is "sufficient evidence" of carcinogenicity for 98 of the 190 chemicals evaluated in its bioassay program,¹⁴⁴ some of which are the same as those reviewed by the International Agency for Research on Cancer.

As stated earlier, sufficient evidence exists for 18 chemicals that they are carcinogenic in humans.⁴⁰

For most of the 200-odd animal carcinogens for which there is "sufficient evidence," it is unlikely that we will ever know with certainty whether they cause cancer in humans because of the difficulty in obtaining appropriate populations suitable for epidemiologic studies. Since it is unlikely we will ever confirm or deny the apparent carcinogenic potential of these 200 chemicals, it appears prudent in the interim to control exposures to them as if they had demonstrated effects in humans.

Short-term Tests for Mutagenicity, DNA Damage and Cell Transformation

Short-term tests¹⁵⁷⁻¹⁵⁹ generally evaluate the ability of a substance to produce mutations, chromosomal alterations or DNA damage in a test organism, or to induce transformation of cultured mammalian cells. Systems that are used in short-term tests include microorganisms (such as bacteria, yeast and molds), cultured mammalian cells and whole animals. These tests are comparatively inexpensive (\$5,000 to \$10,000 per battery of short-term tests versus \$500,000 for an animal bioassay) and can be completed in a relatively short time. A number of these tests can be done, therefore, with limited resources. They offer the potential for providing useful information on the two most intransigent problems in carcinogenic risk assessment—species differences between rodents and humans, and estimating risks at very low doses. Some short-term tests can be performed with human cells or tissues; and some effects, such as DNA damage, can be measured at very low doses.

Use of short-term tests to predict carcinogenicity is justified on both theoretical and empirical grounds. Many of these tests detect biological activities (mutagenicity and cell transformation) that are believed to be stages of carcinogenesis. Most known animal and human carcinogens have been shown to be mutagenic when tested in a suitable battery of short-term tests, while most noncarcinogens have not.¹⁶⁰⁻¹⁶⁹ Because of this, a battery of tests can be a useful predictor of carcinogenicity. Such a strong correlation may exist because most of the chemical carcinogens tested thus far act by mechanisms that involve DNA damage, though this has not been rigorously proved. The particular relevance of tests that measure cell transformation is based on the observation that transformed cells, when implanted into a receptive animal host (such as a "nude" mouse) will form malignant tumors.

There is a high probability that a chemical that is *positive* in an appropriate battery of short-term tests will prove to be a carcinogen when adequately tested in animal cancer tests. Short-term tests can be used, therefore, to augment evidence for carcinogenicity from animal cancer bioassays that, for some reason, are not by themselves definitive. Short-term tests can also indicate the *potential* for carcinogenic hazard of chemicals not yet tested in animals. At present, short-term tests are not sufficiently standardized and validated to

provide *definitive* information about carcinogenicity or noncarcinogenicity in the absence of other evidence.

Risk Assessment and Hazard Evaluation

Quantitative estimates of cancer risks in humans based on extrapolation from animal data are difficult to interpret but are routinely made. What information needs to be developed to permit meaningful estimates of cancer risks from exposures to combinations of chemicals such as occurs in occupational settings? What is a reasonable occupational health policy in the interim? Some principles are emerging. Although differences may exist between species in host responsiveness (differences in pharmacokinetics and DNA-repair efficiencies, for instance), the carcinogenic potencies of chemicals in different species of test animals (rats and mice) are generally similar.¹⁷⁰ Moreover, the responsiveness of test animals is reasonably similar to that of humans for those chemicals (21) that have been examined.^{170,171} Clearly, the inadequacy of human exposure data limits the accuracy of such comparisons. Such *interspecies* differences as exist between rodents and humans must be viewed in relation to the presumed large variation among individuals (such as genetic heterogeneity and host-response differences) in the human population.

Information is needed to improve the accuracy of quantitative estimates of cancer risk from exposures to combinations of agents. In the interim, carcinogenic risks to humans exposed to single or combinations of chemicals must be estimated by extrapolation from available bioassay data using suitable (such as a multi-stage model) methods, while acknowledging that such methods will *underestimate* the true risks if synergisms occur. Compared with the population of test animals, the population of workers is genetically diverse and is simultaneously exposed to a large number of chemicals. Suitable corrections for these differences also should be made in any appropriate risk calculations.

Summaries of hazard evaluations carried out by HESIS for several substances are given below. The evidence for carcinogenicity or reproductive toxicity of these chemicals is based largely on results from studies in test animals.

Ethylene Oxide

Ethylene oxide¹⁰ is a gas that is familiar to most health professionals as a widely used sterilant for heat-sensitive hospital supplies and equipment.¹⁰ Recent studies have indicated that it is carcinogenic by inhalation in male and female rats (leukemia) at dose levels below the current permissible exposure level (50 ppm).^{10,172} It is also mutagenic in 13 test systems,¹⁰ including mammalian somatic cells (micronucleus test)¹⁷³ and germ cells (dominant lethal^{174,175} and heritable translocation tests).¹⁷⁵ In addition, increased frequencies of sister-chromatid exchanges are found in the chromosomes of workers exposed to ethylene oxide¹⁷⁶ at doses less than one-tenth of the current permissible exposure level.¹⁷⁷ Similar chromosomal effects are produced in test animals (monkeys and rabbits).¹⁷⁸ Adverse reproductive

effects (reductions in litter size of exposed female rats at 100 ppm,¹⁷⁹ and sperm count in monkeys at 50 ppm)¹⁰ are evident in test animals at dose levels close to the permissible exposure level.

There is inadequate epidemiologic evidence to assess the carcinogenic or adverse reproductive effects of ethylene oxide in humans.¹⁸⁰⁻¹⁸³ However, because of the extensive evidence for genotoxicity in test animals, HESIS issued a Hazard Alert²³ on this chemical and recommended that the permissible exposure level be reduced and that extensive programs in equipment maintenance, environmental monitoring, training and education, and medical surveillance be initiated in hospitals using this substance.

Hospitals are not normally considered to be workplaces where exposures to toxic substances occur. However, the example of ethylene oxide indicates that members of the medical community, like workers in other occupational settings, need to be aware of the potential health hazards of toxic substances present in their work environment.

Formaldehyde

Like ethylene oxide, formaldehyde is a gas for which there is extensive evidence of genotoxicity and carcinogenicity. Formaldehyde is carcinogenic in two strains of rats and in one strain of mice (nasal cavity squamous cell carcinoma)^{184,185} (Kern WD, Paukov KL, Donofri DJ, et al: Inhalation carcinogenicity of chronic formaldehyde exposure in rats and mice, unpublished data, 1982) at dose levels (6 to 15 ppm) near the current permissible exposure level (2 ppm). It causes DNA damage, mutation or transformation in four test species (bacteria,^{186,187} yeast,¹⁸⁸ *Drosophila melanogaster*¹⁸⁹ and mammalian cells¹⁹⁰) as well as chromosome aberrations in four species.¹⁹¹ Epidemiologic evidence available at present has not shown a carcinogenic effect of formaldehyde in humans.¹⁹²⁻¹⁹⁴ (Fayerweather WE, Pell S, Bender JR: Case-control study of cancer deaths in Du Pont workers with potential exposure to formaldehyde, unpublished report from E. I. Du Pont de Nemours & Co., May, 1982.) However, many of these studies had limited power to detect a significant increase in cancer incidence, and little or no information on exposure levels of formaldehyde was available.

Because the genetic toxicology of formaldehyde is very similar to that of ethylene oxide, it seems reasonable to adopt similarly safe work practices for formaldehyde and extend the health consciousness of the medical community into other work environments. A significant difference between the two chemicals is that formaldehyde is a potent nasal irritant and, thus, has good warning properties whereas ethylene oxide is a nearly odorless, nonirritating substance and, consequently, has poor warning properties. (In fact, a major health hazard from formaldehyde exposure may be respiratory sensitization.)

Ethylene Dibromide

Ethylene dibromide (EDB)^{19,20} has major uses as a lead scavenger in leaded gasoline and as a soil fumigant

in agriculture. With the Med-fly crisis in the summer of 1981, a major increase in use of the chemical was proposed to fumigate citrus fruits, cherries and plums as a means of limiting the spread of Med-fly larvae.

HESIS staff had previously targeted ethylene dibromide as a priority chemical because its genetic toxicology was strikingly similar to that of the closely related brominated compound, DBCP.^{19,20} Both chemicals are potent carcinogens, mutagens and spermatotoxins with effects occurring in test animals at dose levels close to the permitted exposure levels in workers²⁰ (see Table 5). They are among the most potent carcinogens of the widely used industrial chemicals that have been tested in animal bioassays¹⁹² (see Table 6). Ethylene dibromide and DBCP are carcinogenic in rats and mice of both sexes, producing the same malignant tumors at the tissue of first contact (oral route: stomach squamous cell carcinomas;^{3,196} inhalation: respiratory tract carcinomas^{197,198}). For ethylene dibromide, these effects occurred between 1 and 10 mg per kg of body weight per day, uncomfortably close to the dose that workers could

receive (7 mg per kg of body weight per day)¹⁹² at the then current permissible exposure level (20 ppm).

As with DBCP, the chemical is also a potent mutagen in several short-term tests for mutagenicity, including the Ames *Salmonella* test,¹⁹⁹ and *Drosophila*.²⁰⁰ Also, like DBCP, it is a proved animal testicular toxin, causing effects on sperm or male-transmitted reproductive outcomes in three species where it is roughly as potent as DBCP (although in different species and by different routes of exposure, see Table 5).²⁰ To date, there have been no published reports of infertility or induced semen defects in humans associated with exposure to ethylene dibromide.^{207,208}

Because of the extensive evidence that ethylene dibromide is a potent carcinogen and testicular toxin in test animals, HESIS issued a Hazard Alert²⁵ and recommended that the permissible exposure level be reduced (it is now 130 ppb in California). As was the case with ethylene oxide²³ and with the glycol ethers²⁴ (see below), the Hazard Alert was distributed to all companies, unions and workers in California who had potential exposures to the chemical. It is to be hoped that the lessons of DBCP have been learned so that human disease from exposure to these chemicals need never be observed.

Polychlorinated Biphenyls

Polychlorinated biphenyls (PCB's)¹⁴ have received considerable public notoriety as ubiquitous contaminants of our environment. Because they are complex mixtures that are chemically inert, heat stable and non-flammable, they have a variety of uses as dielectric fluids in transformers, hydraulic lubrication fluids, plasticizers and as components in inks, paints and adhesives.¹⁴ They are carcinogenic in male mice (liver carcinomas)²¹⁰ and in two strains of female rats.^{209,211} They cause transformation of mammalian cells in culture (DH Norback, PhD, and R Weltman, PhD, University of Wisconsin Medical School, oral communication, 1980). However, polychlorinated biphenyls are only 1/100 as potent carcinogenic agents in test animals as ethylene oxide, formaldehyde, ethylene dibromide and DBCP, and the evidence for their genotoxicity is less extensive (see Table 6).

The highly chlorinated isomers of polychlorinated

TABLE 5.—Effects of Ethylene Dibromide (EDB) and 1,2-Dibromo-3-chloropropane (DBCP)

	Route	Dose	Effect
Potent sperm toxins			
DBCP:			
Rats ^a	Inhalation	1.2	↓Atrophy
Rabbits ²⁰¹	Inhalation	0.5	↓Count
EDB:			
Cattle ²⁰²⁻²⁰⁴	Oral	2	↓All
Rams ²⁰⁵	Subcutaneous	8-14	↓Morphology
Rats ²⁰⁶	Injection	10	↓Litter size
Species	Oral	Inhalation	
Tumors produced from oral and inhalation exposures to DBCP and EDB			
Mice ♂, ♀	Stomach squamous cell carcinomas	Nasal cavity adenocarcinomas	
Rats ♂, ♀	Stomach squamous cell carcinomas	Respiratory tract carcinomas	

TABLE 6.—Chronic Toxicity of Widely Used Industrial Chemicals

Chemical	1981 US Production (in millions of lbs.)	Use	Carcinogenicity		Relative Dose to Produce Response*	Mutagenicity (Number of Different Short-Term Test Systems)	Permissible Exposure Level (PEL) (ppb)
			Rats	Mice			
1,2-Dibromo-3-chloropropane (DBCP)	Not produced	Fumigant	4+	4+	1	+(5)	2
Ethylene dibromide (EDB)	250	Gasoline additive; fumigant	4+	4+	1	+(3)	125
Ethylene oxide (EtO)	5,000	Intermediate; sterilant	3+	+	10	+(13)	2,000
Formaldehyde (HCHO)	6,000	Intermediate	2+	+	10	+(7)	2,000
Polychlorinated biphenyls (PCB's)	Not produced	Dielectric		3+	100	Cell transformation (1)	
Trichloroethylene (TCE)	130 (1977)	Solvent	+	4+	1,000	+(4)	25,000
Glycol ethers (GE)	1,225	Solvent			...	-(7)	

*Relative dose (mg/kg/day) required to produce significant tumors (DBCP = 1; approximate values).

biphenyls are not readily metabolized in mammalian species, including humans, and tend to bioaccumulate in adipose tissues.¹⁴ Thus, *although the carcinogenic risk is less well established for these chemicals than for the other chemicals already discussed, it is amplified by the long residence time of these agents in the body.* Because of these factors, HESIS recommended¹⁴ that exposure to polychlorinated biphenyls be kept to a minimum.

Glycol Ethers

Glycol ethers¹¹ are a group of widely used industrial solvents found in cleaning and thinning agents and in coatings such as epoxies, wood stains, varnishes, paint and ink. They also serve as surfactants in various extraction processes and as fixatives in perfumes, cosmetics and soaps.¹¹

Recent studies have shown that the glycol ethers and derivatives produce adverse reproductive effects in laboratory animals,¹¹ although they are negative in a variety of short-term tests for mutagenicity.^{212,213} They induced birth defects in the offspring of exposed females and caused retarded sperm development and testicular growth in exposed males in three species (mice,^{214,215} rats^{215,216} and rabbits^{215,216}). These effects occurred by a variety of dose routes (dermal, ingestion and inhalation) and at dose levels that were only slightly higher than the permissible exposure level for these compounds. The congenital defects observed were severe, including neural tube defects in mice (spina bifida and exencephaly)²¹⁴ and cardiac abnormalities in rabbits.^{216,217} Rabbits were particularly sensitive to the testicular effects of ethylene glycol monomethyl ether, and degenerative changes in the testicular germinal epithelium were evident in one study from exposure to 30 ppm for 90 days (permissible exposure level: 25 ppm)²¹⁸ (Miller RR, Calhoun LL, Yano BL: Ethylene glycol monomethyl ether: 13-week vapor inhalation study with male rabbits, unpublished report from Dow Chemical, USA, submitted by the Chemical Manufacturers Association, 1982). Moreover, cardiovascular defects were observed in the offspring of rabbits that received ethylene glycol monoethyl ether.

There is inadequate evidence to assess the reproductive effects of glycol ethers in humans. Women exposed to a variety of solvents, including a glycol ether, reported a variety of gynecological disorders, and their newborn children had a higher incidence of birth defects (including congenital heart defects) than did newborns from a control group.²¹⁹ The significance of the birth abnormalities is tempered by the workers' exposures to other solvents.

Because of the severity of the reproductive effects in test animals, HESIS issued a Hazard Alert²⁴ and recommended that the permissible exposure level for the glycol ethers be lowered and that work practices be changed (use of impermeable gloves, protective clothing and good ventilation) to reduce skin and inhalation exposures. It is noteworthy that male and female reproductive tissues are affected by such nonreactive, nongenotoxic substances as the glycol ethers. Other

solvents that share the special hydrophilic and lipophilic solvating ability of these substances could produce similar effects and should be tested.

Conclusions

Research on cancer and reproductive effects is continuing, and as our understanding increases, programs for prevention and control of these diseases will likely change. Prevention, however, can proceed without precise answers, and we must make occupational health decisions based upon the best available evidence. Cancer risks to the working population and to the individual person can be reduced by a comprehensive program to modify the major identified determinants of cancer and to control exposures to specific carcinogenic substances.²²⁰ Similar programs should be possible with reproductive agents. Modification of work practices cannot be constrained to require absolute certainty when the consequences of inaction could result in serious effects on the health and welfare of the working population.

REFERENCES

1. Whorton D, Krauss RM, Marshall S, et al: Infertility in male pesticide workers. *Lancet* 1977 Dec 17; 2:1259-1261
2. Whorton D, Milby TH, Krauss RM, et al: Testicular function in DBCP exposed pesticide workers. *JOM* 1979 Mar; 21:161-166
3. Bioassay of Dibromochloropropane for possible carcinogenicity. US Dept. of Health, Education, and Welfare publication No. (NIH) 78-828. National Cancer Institute, 1978.
4. Biles RW, Connor TH, Trieff NM, et al: The influence of contaminants on the mutagenic activity of dibromochloropropane (DBCP). *J Environ Pathol Toxicol* 1978 Nov-Dec; 2:301-312
5. Blum A, Ames B: Flame-retardant additives as possible cancer hazards. *Science* 1977; 195:17-23
6. Rosenkranz H: Genetic activity of 1,2-dibromo-3-chloropropane, a widely-used fumigant. *Bull Environ Contam Toxicol* 1975; 14(1):8-12
7. Teramoto S, Saito R, Aoyama H, et al: Dominant lethal mutation induced in male rats by 1,2-dibromo-3-chloropropane (DBCP). *Mutat Res* 1980; 77:71-78
8. Lee IP, Suzuki K: Induction of unscheduled DNA synthesis in mouse germ cells following 1,2-dibromo-3-chloropropane (DBCP) exposure. *Mutat Res* 1979; 68:169-173
9. Torkelson TR, Sadek SE, Rowe VK, et al: Toxicologic investigations of 1,2-dibromo-3-chloropropane. *Toxicol Appl Pharmacol* 1961; 3:545
10. HESIS (Hazard Evaluation System and Information Service): The Toxicity of Ethylene Oxide, With Emphasis on Carcinogenic, Reproductive and Genetic Effects. Berkeley, California Dept. of Health Services, May 1982
11. HESIS (Hazard Evaluation System and Information Service): The Toxicity of Glycol Ethers, With Emphasis on Genetic and Reproductive Effects. Berkeley, California Dept. of Health Services, May 1982
12. HESIS (Hazard Evaluation System and Information Service): An Assessment of the Teratogenic Potential of the Fungicide Captan. Berkeley, California Dept. of Health Services, May 1982
13. HESIS (Hazard Evaluation System and Information Service): Review of the Carcinogenicity, Mutagenicity and Teratogenicity of the Herbicide Nitrofen. Berkeley, California Dept. of Health Services, April 1982
14. HESIS (Hazard Evaluation System and Information Service): The Toxicology of PCBs, An Overview With Emphasis on Human Health Effects and Occupational Exposures. Berkeley, California Dept. of Health Services, January 1981 (updated February 1982)
15. HESIS (Hazard Evaluation System and Information Service): Trichloroethylene (TCE), Evaluation of the Human Health Hazards. Berkeley, California Dept. of Health Services, April 1980 (updated February 1982)
16. HESIS (Hazard Evaluation System and Information Service): Toxicology of Epichlorohydrin. Berkeley, California Dept. of Health Services, February 1982
17. HESIS (Hazard Evaluation System and Information Service): Assessment of the Embryotoxic, Teratogenic and Reproductive Effects of the Fungicide Bayleton® (Triadimefon). Berkeley, California Dept. of Health Services, January 1982
18. HESIS (Hazard Evaluation System and Information Service): Brief Review of the Toxicology of Diazinon. Berkeley, California Dept. of Health Services, October 1981
19. HESIS (Hazard Evaluation System and Information Service): Rationale for an Evaluation of Spermatotoxic Effects in Workers Exposed to Ethylene Dibromide in California. Berkeley, California Dept. of Health Services, September 9, 1981
20. HESIS (Hazard Evaluation System and Information Service): Rationale for a Hazard Alert and a Lowered Exposure Limit for 1, 2-dibromoethane (EDB): A Comparison of the Carcinogenic and Spermatotoxic Effects of EDB and DBCP. Berkeley, California Dept. of Health Services, July 8, 1981

HAZARD EVALUATION SYSTEM

21. HESIS (Hazard Evaluation System and Information Service): Final Report on Potential Health Hazards Associated With the Use of Plastic Pipe in Potable Water Systems. Sacramento, California Dept. of Health Services (Available from California Commission on Housing and Community Development) October 17, 1980
22. HESIS (Hazard Evaluation System and Information Service): 2,4-Dichlorophenoxyacetic Acid (2,4-D), Evaluation of the Human Health Hazards. Berkeley, California Dept. of Health Services, June 16, 1980
23. HESIS (Hazard Evaluation System and Information Service): Hazard Alert on Ethylene Oxide (EtO). Berkeley, California Dept. of Health Services, May 1982 (also available in Spanish)
24. HESIS (Hazard Evaluation System and Information Service): Hazard Alert on Glycol Ethers (Cellosolve® Solvents). Berkeley, California Dept. of Health Services, May 1982 (also available in Spanish)
25. HESIS (Hazard Evaluation System and Information Service): Hazard Alert on Ethylene Dibromide (EDB). Berkeley, California Dept. of Health Services, July 1981 (also available in Spanish)
26. HESIS (Hazard Evaluation System and Information Service): Information Bulletin on Methyl Bromide. Berkeley, California Dept. of Health Services, September 1981 (also available in Spanish)
27. HESIS (Hazard Evaluation System and Information Service): Fact Sheet for Physicians on Arsenic. Berkeley, California Dept. of Health Services, September 1982
28. HESIS (Hazard Evaluation System and Information Service): Fact Sheet for Physicians on Carbon Monoxide. Berkeley, California Dept. of Health Services, September 1982
29. HESIS (Hazard Evaluation System and Information Service): Fact Sheet for Physicians on Cyanide. Berkeley, California Dept. of Health Services, September 1982
30. HESIS (Hazard Evaluation System and Information Service): Fact Sheet for Physicians on Cholinesterase Inhibitor Pesticides: Organophosphates, Carbamates. Berkeley, California Dept. of Health Services, September 1982
31. HESIS (Hazard Evaluation System and Information Service): Fact Sheet for Physicians on Hydrogen Sulfide. Berkeley, California Dept. of Health Services, September 1982
32. HESIS (Hazard Evaluation System and Information Service): Fact Sheet for Physicians on Mercury. Berkeley, California Dept. of Health Services, September 1982
33. HESIS (Hazard Evaluation System and Information Service): Fact Sheet for Physicians on Long-Term Health Effects of Organophosphate Pesticides. Berkeley, California Dept. of Health Services, September 1982
34. Davies DL, Magee BH: Cancer and industrial chemical production. *Science* 1979; 206:1356-1358
35. Doll R, Peto R: Mortality in relation to smoking: 20 years' observation on male British doctors. *Br Med J* 1976; 2:1525-1536
36. Cairns J: Cancer: Science and Society. San Francisco, W. H. Freeman Co, 1978
37. Doll R, Peto R: The causes of cancer: Quantitative estimates of avoidable risks of cancer in the United States today. *JNCI* 1981 Jun; 66:1191-1308
38. The Health Consequences of Smoking—Cancer: A Report of the Surgeon General, publication No. 82-50179. US Dept of Health and Human Services, Public Health Services, Office of Smoking and Health, 1982
39. Peto R, Schneiderman M (Eds): Quantification of Occupational Cancer—Banbury Report No. 9. Cold Spring Harbor, NY, Cold Spring Harbor Laboratory, 1981
40. Althouse R, Huff J, Tomatis L, et al: Chemicals and industrial processes associated with cancer in humans. IARC Monographs, Volumes 1 to 20. IARC Monogr Eval Carcinog Risk Chem Hum (suppl) 1979 Sep; (1):1-71
41. Wyrobek AJ, Burkhardt G, Francis C, et al: Chemically induced alterations of testicular function in man as measured by semen analysis parameters: A report for the gene-tox program. *Mutat Res*, in press
42. Milby TH, Whorton D: Epidemiological assessment of occupationally related, chemically induced sperm count suppression. *JOM* 1980 Feb; 22:77-82
43. Potashnik G, Yanai-Inbar I, Sacks MI, et al: Suppressive effects of 1,2-dibromo-3-chloropropane on human spermatogenesis. *Fertil Steril* 1978 Oct; 30:444-447
44. Egnatz DG, Ott MG, Townsend JC, et al: DBCP and testicular effects in chemical workers: An epidemiological survey in Midland, Michigan. *JOM* 1980 Nov; 22:727-732
45. Lancranian I, Popescu HI, GAvanescu O, et al: Reproductive ability of workmen occupationally exposed to lead. *Arch Environ Health* 1975 Aug; 30:396-401
46. Lancranian I: Alterations of spermatid liquid in patients chronically poisoned by carbon disulfide. *Med Lav* 1972 Jan-Feb; 63:29-33
47. Meyer CR: Semen quality in workers exposed to carbon disulfide compared to a control group from the same plant. *JOM* 1981 Jun; 23:435-439
48. Whorton MD, Milby TH, Stubbs HA, et al: Testicular function among carbaryl-exposed employees. *J Toxicol Environ Health* 1979 Sep; 5:929-941
49. Wyrobek AJ, Watchmaker G, Gordon L, et al: Sperm shape abnormalities in carbaryl-exposed employees. *Environ Health Perspect* 1981; 40:255-265
50. National Institute of Occupational Safety and Health (NIOSH): Reproductive abnormalities in male chemical workers—Kentucky. *Morbidity Mortality Weekly Rep* 1981; 30(17):199
51. Rosenman KD, Anderson HA, Selikoff IJ, et al: Spermatogenesis in men exposed to polybrominated biphenyl (PBB). *Fertil Steril* 1979 Aug; 32:209-213
52. Whorton MD, Stubbs HA, Obrinsky A, et al: Testicular function of men occupationally exposed to para-tertiary butylbenzoic acid. *Scand J Work Environ Health* 1981; 7:204-213
53. Council on Environmental Quality: Chemical Hazards to Human Reproduction. Washington, DC, January 1981
54. Hemminki K, Franssila E, Vainio H: Spontaneous abortions among female chemical workers in Finland. *Int Arch Occup Environ Health* 1980; 45:123-126
55. Funes-Cravioto F, Zapata-Gayon C, Kolmodin-Hedman B, et al: Chromosome aberrations and sister-chromatid exchange in workers in chemical laboratories and a rototyping factory and in children of women laboratory workers. *Lancet* 1977 Aug 13; 2:322-325
56. Meirik O, Kallen B, Gauffin U, et al: Major malformations in infants born of women who worked in laboratories while pregnant (Letter). *Lancet* 1979; 2:91
57. Nordstrom S, Beckman L, Nordenson I: Occupational and environmental risks in and around a smelter in Northern Sweden: I. Variations in birth weight. *Hereditas* 1978; 88:43-46
58. Nordstrom S, Beckman L, Nordenson I: Occupational and environmental risks in and around a smelter in Northern Sweden: III. Frequencies of spontaneous abortion. *Hereditas* 1978; 88:51-54
59. Nordstrom S, Beckman L, Nordenson I: Occupational and environmental risks in and around a smelter in Northern Sweden: V. Spontaneous abortion among female employees and decreased birth weight in their offspring. *Hereditas* 1979; 90:291-296
60. Nordstrom S, Beckman L, Nordenson I: Occupational and environmental risks in and around a smelter in Northern Sweden: VI. Congenital malformations. *Hereditas* 1979; 90:297-302
61. Panova S, Ivanova S: Changes in the ovarian function and some functional indices of liver of women in professional contact with metallic mercury. *Akush Ginekol (Mosk)* 1979; 15:136-137
62. Vaisman AI: Working conditions in surgery and their effect on the health of anesthesiologists. *Eksp Khir Anest* 1967 May-Jun; 12:44-49 (Russian)
63. Askrog V, Harvald B: Teratogen effect of inhalation of anesthesia. *Nord Med* 1970 Apr 16; 83:498-500
64. Cohen EN, Bellville JW, Brown BW: Anesthesia, pregnancy and miscarriage. *Anesthesiology* 1971; 35:343-347
65. Cohen EN: Occupational disease among operating room personnel: A national study. *Anesthesiology* 1974; 41:321-340
66. Cohen EN, Brown BW, Bruce DL, et al: A survey of anesthetic health hazards among dentists. *J Am Dent Assoc* 1975; 90:1291-1296
67. Knill-Jones RP, Rodrigues LV, Moir DD, et al: Anaesthetic practice and pregnancy—Controlled survey of women anaesthetists in the United Kingdom. *Lancet* 1972; 1:1326-1328
68. Knill-Jones RP, Newman BJ, Spence AA: Anaesthetic practice and pregnancy. *Lancet* 1975; 2:807-809
69. Pharoah POD, Alberman E, Doyle P, et al: Outcome of pregnancy among women in anaesthetic practice. *Lancet* 1977; 1:34-36
70. Corbett TH, Cornell RG, Endres JL, et al: Birth defects among children of nurse-anesthetists. *Anesthesiology* 1974; 41:341-344
71. Aldyreva MV, Klimova TS, Izyumova AS, et al: The influence of phthalate plasticisers on the generative function. *Gig Tr Prof Zabol* 1975; 19:25-29
72. Shumilina AV: Menstrual and child-bearing functions of female workers occupationally exposed to the effects of formaldehyde. *Gig Tr Prof Zabol* 1975 Dec; 19:18-21
73. Wiley FH, Hueper WC, Von Oettingen WF: On toxic effects of low concentrations of carbon disulfide. *J Indust Hyg Toxicol* 1936; 18:733-740
74. Ehrhardt W: Experiences with the employment of women exposed to carbon disulfide. In: *International Symposium on Toxicology of Carbon Disulfide*, Prague, 1966. Amsterdam, Excerpta Medica Foundation, 1967, pp 240-244
75. Blekherman NA, Ilyina VI: Changes of ovary function in women in contact with organochlorine compounds. *Pediatrriia* 1973; 52:57-59
76. Veis VP: Some data on the status of the sexual sphere in women who have been in contact with organochlorine compounds. *Pediatr Akush Ginek* 1970; 32:48-49
77. Mukhametova GM, Vozovaya MA: Reproductive power and incidence of gynecological affections among female workers exposed to a combined effect of gasoline and chlorinated hydrocarbons. *Gig Tr Prof Zabol* 1972; 16:6-9
78. Criteria for a Recommended Standard—Occupational Exposure to Toluene Diisocyanate, DHEW publication No. (NIOSH) 73-11023. National Institute for Occupational Safety and Health (NIOSH), 1973
79. Espir MLE, Hall JW, Shirreffs JG, et al: Impotence in farm workers using toxic chemicals. *Br Med J* 1970; 1:423-425
80. Yoder J, Watson M, Benson WW: Lymphocyte chromosome analysis of agricultural workers during extensive occupational exposure to pesticides. *Mutat Res* 1973; 21:335-340
81. Rabello MN, Becak W, DeAlmeida WF, et al: Cytogenetic study on individuals occupationally exposed to DDT. *Mutat Res* 1975; 28:449-454
82. Shabatal F, Bichacho S, Halbrecht I: Cytogenetic observations in infertile men working with insecticidal compounds. *Acta Genet Med Gemellol* 1978; 27:51-56
83. Kiraly J, Szentesi I, Ruzicska M, et al: Chromosome studies in workers producing organophosphate insecticides. *Arch Environ Contam Toxicol* 1979; 8:309-319
84. Sandifer SH, Wilkins RT, Loadholt CB, et al: Spermatogenesis in agricultural workers exposed to dibromochloropropane (DBCP). *Bull Environ Contam Toxicol* 1979; 23:703-710
85. Kapp, RW, Picciano DJ, Jacobson CB: Y-chromosomal nondisjunction in dibromochloropropane-exposed workmen. *Mutat Res* 1979; 64:47-51
86. Nogaki K: On the action of lead on the body of lead refinery workers: Particularly on the conception, pregnancy and parturition in the case of females and on the vitality of their newborn. *Excerpt Med* 1958; (XVII)4:2176 (as reported by USEPA 1977)
87. Fahim MS, Fahim Z, Hall DG: Effects of subtoxic lead levels on pregnant women in the state of Missouri. *Res Commun Chem Pathol Pharmacol* 1976; 13:309-331
88. Schwanitz G, Lehnert G, Gebhart E: Chromosomal injury due to occupational lead poisoning. *Ger Med Month* 1970; 15:738-746

HAZARD EVALUATION SYSTEM

89. Purchase IFH, Richardson CR, Anderson D, et al: Chromosomal analyses in vinyl chloride-exposed workers. *Mutat Res* 1978 Jul; 57:325-334
90. Ducatman A, Hirschhorn K, Selikoff IJ: Vinyl chloride exposure and human chromosome aberrations. *Mutat Res* 1975; 31:163-168
91. Heath CW, Dumont CR, Gamble J, et al: Chromosomal damage in men occupationally exposed to vinyl chloride monomer and other chemicals. *Environ Res* 1977; 14:68-72
92. Funes-Cravioto F, Lambert B, Lindsten J, et al: Chromosome aberrations in workers exposed to vinyl chloride. *Lancet* 1975; 1:459
93. Hansteen I, Hillstad L, Thiiis-Evensen E, et al: Effects of vinyl chloride in man: A cytogenetic follow-up study. *Mutat Res* 1978; 51:271-278
94. Szentesi I, Hornyák E, Ungváry G, et al: High rate of chromosomal aberration in PVC workers. *Mutat Res* 1976 Nov; 37(2-3):313-316
95. Fleig I, Thiess AM: Mutagenicity of vinyl chloride—External chromosome studies on persons with and without VC illness, and on VC exposed animals. *JOM* 1978 Aug; 20:557-561
96. Taylor JR, Selhorst JB, Houff SA, et al: Chlorocone intoxication in man: I: Clinical observations. *Neurology* 1978; 28:626-630
97. Wong O, Utidjian MD, Karten VS: Retrospective evaluation of reproductive performance of workers exposed to ethylene dibromide (EDB). *JOM* 1979; 21:98-109
98. Wyrobek AJ, Gordon LA, Burkhart JG, et al: An evaluation of the mouse sperm morphology assay and sperm assays in other non-human mammals: A report for the gene-tox program. *Mutat Res*, in press
99. Shepard TH: Catalog of Teratogenic Agents. Baltimore and London, The Johns Hopkins University Press, 1980
100. Wilson JG, Fraser FC (Eds): Handbook of Teratology, Comparative, Maternal, and Epidemiologic Aspects. New York and London, Plenum Press, 1977
101. Klingberg MA, Weatherall JA (Eds): Epidemiologic Methods for Detection of Teratogens. Basel, Switzerland, S. Karger, 1979
102. Hatch M, Kline J, Stein Z: Power considerations in studies of reproductive effects of vinyl chloride and some structural analogs. *Environ Health Perspect* 1981; 41:195-201
103. Schardein JL (Ed): Drugs as Teratogens. Cleveland, CRC Press, 1976
104. Kuratsune M, Yoshimura T, Matsuzaha J, et al: Epidemiological study of yusho, a poisoning caused by ingestion of rice oil contaminated with a commercial brand of polychlorinated biphenyls. *Environ Health Perspect* 1972; 1:119-128
105. Allan JR, Barsotti DA: The effects of transplacental and mammary movement of PCBs on infant rhesus monkeys. *Toxicology* 1976 Nov-Dec; 6:331-340
106. Hanson JW, Streissguth AP, Smith DW: The effects of moderate alcohol consumption during pregnancy on fetal growth and morphogenesis. *J Pediatr* 1978 Mar; 92:457-460
107. Sander S, Amels D: The action of aethanol on the prae-natal development of albino rats—An attempt of multiphasic screening. *Rev Roum Embryol Cytol Ser Embryol* 1971; 8:105-118
108. Emerson DJ: Congenital malformation due to attempted abortion with aminopterin. *Am J Obstet Gynecol* 1962; 84:356-357
109. Baranov VS: Characteristics of the teratogenic effect of aminopterin compared with that of other teratogenic agents. *Bull Exp Biol* 1966; 1:77-81
110. Milunsky A, Graef JW, Gaynor MF: Methotrexate-induced congenital malformations. *J Pediatr* 1968; 72:790-795
111. Wilson JG: Use of rhesus monkeys in teratological studies. *Fed Proc* 1971; 30:104-109
112. Amin-Zaki L, Elhassani S, Majeed MA, et al: Perinatal methylmercury poisoning in Iraq. *Am J Dis Child* 1976; 130:1070-1076
113. Khara KS: Teratogenic effects of methylmercury in the cat: Note on the use of this species as a model for teratogenicity studies. *Teratology* 1973; 8:293-304
114. Khara KS, Tabacova SA: Effects of methylmercuric chloride on the progeny of mice and rats treated before or during gestation. *Food Cosmetics Toxicol* 1973; 11:245-254
115. Sex hormones (II). IARC Monogr Eval Carcinog Risk Chem Hum 1979 Dec; 21:1-561
116. Hendrickx AG, Benirschke K, Thompson RS, et al: The effects of prenatal diethylstilbestrol (DES) exposure on the genitalia of pubertal Macaca mulatta—I. Female offspring. *J Reprod Med* 1979 May; 22:233-240
117. Hanson JW, Smith DW: The fetal hydatoid syndrome. *J Pediatr* 1975; 87:285-290
118. Harbison RD, Becker BA: Relation of dosage and time of administration of diphenylhydantoin to its teratogenic effect in mice. *Teratology* 1969; 2:305-312
119. Hammond EC, Selikoff IJ, Seidman H: Asbestos exposure, cigarette smoking and death rates. In *Health hazards of asbestos exposure*. Ann NY Acad Sci 1979; 330:473-490
120. National Research Council: Diet, Nutrition, and Cancer. Washington, DC, National Academy Press, 1982
121. World Health Organization (WHO) Scientific Group: Principles for the Testing and Evaluation of Drugs for Carcinogenicity. WHO Technical Report, Series No. 426, 1969
122. Food Safety Council: Proposed System for Food Safety Assessment. Final Report of the Scientific Committee, Washington, DC, 1980
123. Safe Drinking Water Committee, Advisory Center on Toxicology, National Research Council, National Academy of Sciences: Drinking Water and Health. Washington, DC, 1977
124. US Occupational Safety and Health Administration (OSHA): Identification, Classification, and Regulation of Potential Occupational Carcinogens, 45FR5001-5296, January 22, 1980
125. US Inter-Agency Regulatory Liaison Group (IRLG): Scientific Basis for Identification of Potential Carcinogens and Estimation of Risk, 44FR39858, July 6, 1979, and JNCI 1979; 63:244-268
126. US Office of Science and Technology Policy (OSTP): Identification, Characterization, and Control of Potential Human Carcinogens: A Framework for Federal Decision-Making. Executive Office of the President, 1979, and JNCI 1980; 64:169-176
127. US Congress Office of Technology Assessment (OTA): Assessment of Technologies for Determining Cancer Risks From the Environment. US Congress, 1981
128. Council on Environmental Quality (CEQ), Toxic Substances Strategy Committee: Toxic Chemicals and Public Protection: A Report to the President, May 1980
129. National Cancer Advisory Board, Subcommittee on Environmental Carcinogenesis: General Principles for Assessing the Evidence for Carcinogenicity of Chemical Substances. JNCI 1977; 58:461-464
130. Doll R: Strategy for detection of cancer hazards to man. *Nature* 1977 Feb 17; 265(5595):589-596
131. Karstadt M: A survey of epidemiologic data on humans exposed to animal carcinogens. In Peto R, Schneiderman M (Eds): Quantification of Occupational Cancer, Banbury Report No. 9. Cold Spring Harbor, NY, Cold Spring Harbor Laboratory, 1981, pp 223-245
132. Day NE, Brown CC: Multistage models and the primary prevention of cancer. JNCI 1980; 64:977-989
133. Doll R, Peto R: Mortality in relation to smoking: 20 years' observation on male British doctors. *Br Med J* 1976 Dec 25; 2:1525-1536
134. Sontag JM, Page NP, Saffioti U: Guidelines for carcinogenic bioassay in small rodents. National Cancer Institute, Bethesda, MD, 1976
135. US Environmental Protection Agency (USEPA): Proposed Health Effects Test Standards for Toxic Substances Control Act Test Rules and Proposed Good Laboratory Practice Standards for Health Effects, 44FR27334-27375, May 9, 1979
136. US Environmental Protection Agency (USEPA): Proposed Health Effects Test Standards for Toxic Substances Control Act Test Rules and Proposed Good Laboratory Practice Standards for Health Effects, 44FR-44054-44093, July 26, 1979
137. US Environmental Protection Agency (USEPA): Support Document Test Data Development Standards: Chronic Health Effects. Toxic Substances Control Act, Section 4, 1979
138. Long-term and short-term screening assays for carcinogens: A critical appraisal. IARC Monogr Eval Carcinog Risk Chem Hum (Suppl) 1980; (2 Suppl):1-426
139. Organization for Economic Cooperation and Development (OECD): OECD Short-term and Long-term Toxicology Groups, Final Report. Paris, December 31, 1979
140. Gart JJ, Chu KC, Tarone RE: Statistical issues in interpretation of chronic bioassay tests for carcinogenicity. JNCI 1979 Apr; 62:957-974
141. Purchase IFH: Inter-species comparisons of carcinogenicity. *Br J Cancer* 1980 Mar; 41:454-468
142. Tomatis L, Partensky C, Montesano R: The predictive value of mouse liver tumour induction in carcinogenicity testing—A literature survey. *Int J Cancer* 1973 Jul 15; 12:1-20
143. Tomatis L, Agthe C, Bartsch H, et al: Evaluation of the carcinogenicity of chemicals: A review of the Monograph Program of the International Agency for Research on Cancer (1971-1977). *Cancer Res* 1978 Apr; 38:877-885
144. Griesemer RA, Cueto C Jr: Toward a classification scheme for degrees of experimental evidence for the carcinogenicity of chemicals for animals. IARC Sci Publ 1980; (27):259-281
145. Chu KC, Cueto C, Ward JM: Factors in the evaluation of 200 National Cancer Institute carcinogen bioassays. *J Toxicol Environ Health* 1981; 8:251-280
146. International Agency for Research on Cancer (IARC): An evaluation of chemicals and industrial processes associated with cancer in humans based on human and animal data: IARC Monographs 1-20. *Cancer Res* 1980; 40:1-12
147. International Agency for Research on Cancer (IARC): IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans, Vols 1-26, Lyon, France, 1972-1982
148. Tomatis L: The predictive value of rodent carcinogenicity tests in the evaluation of human risks. *Annu Rev Pharmacol Toxicol* 1979; 19:511-530
149. Bishop JM: Oncogenes. *Sci Am* 1982 Mar; 246:82-92
150. Marx JL: Cancer cell genes linked to viral oncogenes. *Science* 1982 May; 216:724
151. Marx JL: Gene transfer yields cancer clues. *Science* 1982 Feb; 215:955-957
152. Shih C, Shilo BZ, Goldfarb MP, et al: Passage of phenotypes of chemically transformed cells via transfection of DNA and chromatin. *Proc Natl Acad Sci USA* 1979 Nov; 76:5714-5718
153. Shih C, Padhy LC, Murray M, et al: Transforming genes of carcinomas and neuroblastomas introduced into mouse fibroblasts. *Nature* 1981 Mar; 290:261-264
154. Bishop JM: Enemies within: The genesis of retrovirus oncogenes. *Cell* 1981 Jan; 23:5-6
155. Eva A, Robbins KC, Andersen PR, et al: Cellular gene analogous to retroviral oncogenes are transcribed in human tumour cells. *Nature* 1982 Jan; 295:116-119
156. Goldfarb M, Shimizu K, Peruchio M, et al: Isolation and preliminary characterization of a human transforming gene from T24 bladder carcinoma cells. *Nature* 1982 Apr; 296:404-409
157. Stich HF, San RH (Eds): Short-Term Tests for Chemical Carcinogens. New York, Springer-Verlag, 1981
158. Sugimura T, Kondo S, Takebe H (Eds): Environmental Mutagens and Carcinogens. Tokyo/New York, University of Tokyo Press/Alan R. Liss, 1982
159. Hollstein M, McCann J, Angelosanto FA, et al: Short-term tests for carcinogens and mutagens. *Mutat Res* 1979 Sep; 65:133-226
160. McCann J, Ames BN: The Salmonella/microsome mutagenicity test: Predictive value for animal carcinogenicity. In Hiatt HH, Watson JD, Winsten JA (Eds): Human Cancer—Book C: Human Risk Assessment. Cold Spring Harbor, New York, Cold Spring Harbor Laboratory, 1977, pp 1431-1450

161. McCann J, Ames BN: Detection of carcinogens as mutagens in the *Salmonella/microsome* test: Assay of 300 chemicals. *Proc Natl Acad Sci USA* 1975 Dec; 72:5135-5139
162. McCann J, Ames BN: Detection of carcinogens as mutagens in the *Salmonella/microsome* test: Assay of 300 chemicals: Discussion. *Proc Natl Acad Sci USA* 1976 Mar; 73:950-954
163. Rinkus SJ, Legator MS: Chemical characterization of 465 known or suspected carcinogens and their correlation with mutagenic activity in the *Salmonella typhimurium* system. *Cancer Res* 1979 Sep; 39:3289-3318
164. Ames BN, McCann J: Validation of the *Salmonella* test: A reply to Rinkus and Legator. *Cancer Res* 1981; 41:4192-4196
165. Sugimura T, Scott D, Bridges BA, et al: A critical review of sub-mammalian systems for mutagen detection. In Scott D, Bridges BA, Sobels FH (Eds): *Progress in Genetic Toxicology—Vol 2. Developments in Toxicology and Environmental Science*. Amsterdam, Elsevier/North-Holland, 1977
166. Bridges BA: Short-term tests and human health—The central role of DNA repair. In Emmelot P, Kriek E (Eds): *Environmental Carcinogenesis: Occurrence Risk Evaluation and Mechanisms*. Amsterdam, Elsevier/North-Holland, 1979
167. McMahon RE, Cline JC, Thompson CZ: Assay of 855 test chemicals in ten tester strains using a new modification of the Ames test for bacterial mutagens. *Cancer Res* 1979; 39:682-693
168. Bartsch H, Malaveille C, Camus AM, et al: Bacterial and mammalian mutagenicity tests: Validation and comparative studies on 180 chemicals. *IARC Sci Publ* 1980; (27):179-241
169. de Serres FJ, Shelby MD (Eds): *Comparative Chemical Mutagenesis*. New York, Plenum Press, 1981
170. Crouch E, Wilson R: Interspecies comparison of carcinogenic potency. *J Toxicol Environ Health* 1979 Nov; 5:1095-1118
171. Meselson M: Pest Control: An Assessment of Present and Alternative Technologies—Vol 1: Contemporary Pest Control Practices and Prospects: The Report of the Executive Committee, National Academy of Sciences (NAS). Washington, DC, 1975, pp 66-85
172. Snellings WM, Weil CS, Maronpot RR: Final Report on Ethylene Oxide Two-year Inhalation Study on Rats, Project Report 44-20. Bushy Run Research Center, Union Carbide, January 28, 1981 (Submitted to HESIS by Union Carbide, February 1982.)
173. Appelgren LE, Eneroth G, Grant C, et al: Testing of ethylene oxide for mutagenicity using the micronucleus test in mice and rats. *Acta Pharmacol Toxicol (Copenh)* 1978 Jul; 43:69-71
174. Embree JW, Lyon JP, Hine CH: The mutagenic potential of ethylene oxide using the dominant-lethal assay in rats. *Toxicol Appl Pharmacol* 1977; 40:261-267
175. Generoso WM, Cain KT, Krishna M, et al: Heritable translocation and dominant-lethal mutation induction with ethylene oxide in mice. *Mutat Res* 1980; 73:133-142
176. Garry V, Hozier J, Jacobs D, et al: Ethylene oxide: Evidence of human chromosomal effects. *Environ Mutagen* 1979; 1:375-382
177. Yager JL: Sister-Chromatid Exchange as a Biological Monitor for Workplace Exposure: A Comparative Study of Exposure-Response to Ethylene Oxide. Berkeley, Northern California Occupational Health Center Report No. NCOHC 82-B-2, 1982
178. Yager JW, Benz RD: Sister chromatid exchanges induced by ethylene oxide after inhalation exposure in the rabbit. *Environ Mutagen*, in press
179. Carnegie-Mellon Institute of Research: Final Report on Ethylene Oxide One-Generation Reproduction Inhalation Study, Project Report 42-7. Carnegie-Mellon Institute of Research, May 1, 1979 (Submitted to HESIS by Union Carbide Corporation.)
180. Morgan RW, Claxton KW, Divine BJ, et al: Mortality among ethylene oxide workers. *JOM* 1981; 23:767-770
181. Hogstedt C, Malmqvist N, Wadman B: Leukemia in workers exposed to ethylene oxide. *JAMA* 1979 Mar 16; 241:1132-1133
182. Hogstedt C, Rohlén O, Berndtsson BS, et al: A cohort study of mortality and cancer incidence in ethylene oxide production workers. *Br J Ind Med* 1979 Nov; 36:276-280
183. Abrahams RH: Recent studies with workers exposed to ethylene oxide. In Jorkasky JF (Ed): *The Safe Use of Ethylene Oxide: Proceeding of the Educational Seminar, Health Industry Manufacturers Association*. Washington, DC, HIMA Report No. 80-4, 1980; pp 27-38, 211-220
184. Albert RE, Sellakumar AR, Laskin S, et al: Nasal cancer in the rat induced by gaseous formaldehyde and hydrogen chloride. *JNCI*, in press
185. Kerns WD, Donofrio DJ, Pavkov KL: The chronic effects of formaldehyde vapor inhalation in rats and mice—A preliminary report. In *Proceedings of the Third CIIT Annual Conference, Formaldehyde Toxicity*. New York, Hemisphere Pub Corp, in press
186. Rosenkranz HS, Leifer Z: Determining the DNA-modifying activity of chemicals using DNA-polymerase-deficient *Escherichia coli*. In de Serres FJ, Hollaender A (Eds): *Chemical Mutagens—Vol 6*. New York, Plenum Press, 1980, pp 109-147
187. Sasaki Y, Endo R: Mutagenicity of aldehydes in *Salmonella* (Abstract No. 27). *Mutat Res* 1978; 54:251-252
188. Chanet R, Izard C, Moustacchi E: Genetic effects of formaldehyde in yeast: I. Influence of the growth stages on killing and recombination. *Mutat Res* 1975; 33:179-186
189. Auerbach C, Moutschen-Dahmen M, Moutschen J: Formaldehyde sensory irritation. In *Proceedings of the Third Annual CIIT Conference: Formaldehyde Toxicity*. New York, Hemisphere Pub Corp, in press
190. Ragan DI, Boreiko CJ: Initiation of C3H/10-1/2 cell transformation by formaldehyde. *Cancer Letters* 1981; 13:325-331
191. Manna GK, Parida BB: Formalin-induced sex chromosome breakage in the spermatocyte cells of the grasshopper. *Tristria Pulvinata J Cytol Genet (India)* 1967; 2:86-91
192. Walrath J, Fraumeni JR Jr: Proportionate mortality among New York embalmers. In *Proceedings of the Third Annual CIIT Conference: Formaldehyde Toxicity*. New York, Hemisphere Pub Corp, in press
193. Wong O: An epidemiologic mortality study of a cohort of chemical workers potentially exposed to formaldehyde, with a discussion on SMR and PMR. In *Proceedings of the Third Annual CIIT Conference: Formaldehyde Toxicity*. New York, Hemisphere Pub Corp, in press
194. Marsh GM: Proportional mortality among chemical workers exposed to formaldehyde. In *Proceedings of the Third Annual CIIT Conference, Formaldehyde Toxicity*. New York, Hemisphere Pub Corp, in press
195. Ames BN, Gold LS, Havender WR, et al: Carcinogenic Potency, California Policy Report Seminar Final Report No. 3. Berkeley, University of California Institute of Governmental Studies, December 1981
196. Bioassay of 1,2-Dibromoethane for Possible Carcinogenicity, DHEW publication No. 78-1336. 1978, National Cancer Institute
197. Bioassay of 1,2-Dibromoethane for Possible Carcinogenicity, DHEW publication No. 81-1766. 1981, National Cancer Institute
198. Bioassay of 1,2-Dibromo-3-Chloropropane for Possible Carcinogenicity, DHEW publication No. 82-1762. 1982, National Cancer Institute
199. Rosenkranz HS: Mutagenicity of halogenated alkanes and their derivatives. *Environ Health Perspect* 1977 Dec; 21:79-84
200. Kale P, Baum JW: Sensitivity of *Drosophila melanogaster* to low concentrations of gaseous mutagens: III. Dose-related effects. *Environ Mutag* 1981; 3(1):65-70
201. Burek JD, Murray FJ, Rao KS, et al: Pathogenesis of inhaled 1,2-dibromo-3-chloropropane (DBCP) induced testicular atrophy in rats and rabbits. *Proceedings and abstracts of the 18th annual meeting of the Society of Toxicology*, Boston, 1982
202. Amir D, Volcani R: The effect of dietary ethylene dibromide (EDB) on the testes of bulls. *Fertil Steril* 1967 Jan-Feb; 18:144-147
203. Amir D, Lavon U: Changes in total nitrogen, lipoproteins and amino acids in epididymal and ejaculated spermatozoa of bulls treated orally with ethylene dibromide. *J Reprod Fertil* 1976 May; 47:73-76
204. Amir D, Esnault C, Nicolle JC, et al: DNA and protein changes in the spermatozoa of bulls treated orally with ethylene dibromide. *J Reprod Fertil* 1977 Nov; 51:453-456
205. ElJack AH, Hrudka F: Pattern and dynamics of teratospemia induced in rats by parenteral treatment with ethylene dibromide. *J Ultrastruct Res* 1979 May; 67:124-134
206. Short RD, Winston JM, Hong CB, et al: Effects of ethylene dibromide on reproduction in male and female rats. *Toxicol Appl Pharmacol* 1979 Jun; 49:97-105
207. Wong O, Utidjian HM, Katten VS: Retrospective evaluation of reproductive performance of workers exposed to ethylene dibromide (EDB). *JOM* 1979 Feb; 21:98-102
208. Letz G: Testimony to Cal/OSHA Standards Board hearing on ethylene dibromide, Sacramento, Calif., 1981
209. Kimbrough RD, Linder RE: Induction of adenofibrosis and hepatomas of the liver in BALB/c mice by polychlorinated biphenyls (Aroclor 1254). *JNCI* 1974 Aug; 53:547-552
210. Ito N, Nagasaki H, Makiura S, et al: Histopathological studies on liver tumorigenesis in rats treated with polychlorinated biphenyls. *Gann* 1974 Dec; 65:545-549
211. Kimbrough RD, Squire RA, Linder RE, et al: Induction of liver tumors in Sherman strain female rats by polychlorinated biphenyl (Aroclor 1260). *JNCI* 1975 Dec; 55:1453-1459
212. McGregor DB: Tier II Mutagenicity Screening of 13 NIOSH Priority Compounds—Individual Compound Report, 2-Methoxyethanol, Report No. 20. Prepared by Inveresk Research International Ltd., under NIOSH Contract No. 210-78-0026, November 1, 1980
213. McGregor DB: Tier II Mutagenicity Screening of 13 NIOSH Priority Compounds—Individual Compound Report, Bis(2-methoxyethoxy)-ether, Report No. 34. Prepared by Inveresk Research International Ltd., under NIOSH Contract No. 210-78-0026, July 7, 1981
214. Nagano K, Nakayama E, Oobayashi H, et al: Embryotoxic effects of ethylene glycol monomethyl ether in mice. *Toxicology* 1981; 20:335-343
215. Blair EH: EPA Document Control No. 8EHQ02810384. Submission to EPA by the Dow Chemical Company under Section 8e of the Toxic Substances Control Act, January 25, 1982
216. Andrew FD, Buschbom RL, Cannon WC, et al: Teratologic Assessment of Ethylbenzene and 2-Ethoxyethanol. Report submitted by Battelle Pacific Northwest Laboratories to NIOSH under Contract No. 210-79-0037, 1981
217. Nagano K, Nakayama E, Koyano M, et al: Testicular atrophy of mice induced by ethylene glycol monoalkyl ethers (author's trans), *Eng abstract. Jap Ind Health* 1979 Jan; 21:29-35
218. Miller RR, Ayres JA, Young JT, et al: Ethylene glycol monomethyl ether: A subchronic inhalation study in rats and rabbits, abstracted. *Toxicologist* 1982; 2(1):11(abst. no. 40)
219. Syrobadko ON, Malsheva ZV: Work conditions and their effect on certain specific functions among women who are engaged in the production of enamel-insulated wire (translated from Russian). *Gig Tr Prof Zabol* 1977; 4:25-28
220. Carcinogen Identification Policy: A Statement of Science as a Basis of Policy. Sacramento, State of California Department of Health Services, July 1982